

Theoretical Isochrones with Extinction in the K-BandSungsoo S. Kim¹, Donald F. Figer², Myung Gyoon Lee³, and Seungkyung Oh¹**ABSTRACT**

We calculate theoretical isochrones, in a consistent way, for five filters in the atmospheric window between $1.9\,\mu\text{m}$ and $2.5\,\mu\text{m}$, K , K' , K_s , F205W, and F222M, using the Padova stellar evolutionary models by Girardi et al. Even when displayed in the same Vega magnitude system, the near-infrared colors of the same isochrone can differ by up to 0.18 mag at its bright end, depending on the filter. We present magnitude transformations between K -band filters as a function of color from H & K band filters. Isochrones with extinction at K of up to 6 mag are also presented. We find that care is needed when comparing extinction values that are estimated using different filter sets in the K -band, in particular when comparing those between atmospheric and space filter sets: extinction values for space filters can be in error by up to 0.3 mag. To reduce this error, we introduce an “effective extinction slope” for each filter set and isochrone model, which describes the extinction behaviour of isochrones in the color-magnitude diagram more correctly than the actual extinction law. Our calculation also suggests that the extinction law implied by the observations of Rieke, Rieke, & Paul for wavelengths between H and K bands is better described by a power-law function with an exponent of 1.61, instead of 1.55, which is commonly used with an assumption that the transmission functions of H and K filters are Dirac delta functions.

Subject headings: Hertzsprung-Russell diagram — techniques: photometric — stars: fundamental parameters — infrared: stars

1. INTRODUCTION

Deep near-infrared stellar photometry is an important tool for studies of the structure and stellar population in the Galactic bulge and the disk, where the extinction by dust is quite significant. In the near-infrared ($1 - 5\,\mu\text{m}$), deep ground-based imaging is generally limited to the three

¹Dept. of Astronomy & Space Science, Kyung Hee University, Yongin-shi, Kyungki-do 449-701, Korea; sskim@ap.khu.ac.kr

²Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218; figer@stsci.edu

³Astronomy Program, SEES, Seoul National University, Seoul 151-742, Korea; mglee@astrog.snu.ac.kr

shorter atmospheric windows, J ($1.2\,\mu\text{m}$), H ($1.65\,\mu\text{m}$), and K ($2.2\,\mu\text{m}$). Longward of $2.5\,\mu\text{m}$, thermal emission from a warm telescope, instrument, and sky (for ground-based observations) makes photometry highly inefficient.

The K -band is the most popular among these three bands, because it suffers from the least extinction and provides the widest wavelength baseline when combined with visible bands. Contamination by the thermal emission in the K filter (Johnson et al. 1966) is not negligible, so astronomers have introduced several filters with slightly shorter long wavelength cutoffs, i.e. K' (Wainscoat & Cowie 1992) and K_s (K -short; developed by M. Skrutskie—see the appendix of Persson et al. 1998), that effectively reduce the total background from a warm sky and telescope by a factor of two.

On the other hand, the near-infrared camera on board *Hubble Space Telescope* (*HST*), NICMOS, has a unique filter system, as it is not subject to the effects of atmospheric absorption. The NICMOS F160W filter and the H filter (Johnson et al. 1966) have similar central wavelengths, and the NICMOS F205W and F222M filters transmit wavelengths that fall within the passband of the Johnson K filter.

As enumerated above, there are at least five different broad- and medium-band filters in common usage that cover wavelengths near $2.2\,\mu\text{m}$: three atmospheric filters K , K' , & K_s , and two space filters F205W & F222M. Transmission functions for these five filters are quite different from each other (see Figure 1), but there have been no studies that consistently compare the magnitudes and colors of stars as observed with these five filters as a function of interstellar extinction.

There have been a few studies that compare theoretical isochrones or colors for *some* of the five filters. Girardi et al. (2002) presented theoretical isochrones for K , K_s , and F205W filters among others, but not for K' and F222M. Origlia & Leitherer (2000) provided color transformations between $J - H$ & F110W–F160W, and between $J - K$ & F110W–F222M, and Stephens et al. (2000) presented transformation between K magnitude and F110W–F205W color, but none for K' , K_s , or F205W. Wainscoat & Cowie (1992) investigated the transformation between the K' and K filters.

In the present paper, we present and analyze theoretical isochrones that are calculated in a consistent way for the five filters near the $2\,\mu\text{m}$ atmospheric window over a range of extinctions.

The magnitude system we adopt here is explained in § 2, and the stellar spectral library and evolutionary tracks we use are discussed in § 3. Theoretical isochrones are presented in § 4, and our findings are then summarized in § 5.

2. Magnitude System

Here, we briefly discuss our adopted magnitude system, following the discussion by Girardi et al. (2002).

The flux of a star measured at the top of the Earth’s atmosphere, f_λ , is related to the flux at the stellar photosphere, F_λ , by

$$f_\lambda = 10^{-0.4A_\lambda} (R/d)^2 F_\lambda, \quad (1)$$

where A_λ is the extinction in magnitudes at the wavelength λ , R is the stellar radius, and d is the distance to the star. The apparent magnitude, m_{S_λ} , as measured using a filter with transmission function S_λ is given by

$$m_{S_\lambda} = -2.5 \log \left(\frac{\int_{\lambda_1}^{\lambda_2} f_\lambda S_\lambda d\lambda}{\int_{\lambda_1}^{\lambda_2} f_\lambda^0 S_\lambda d\lambda} \right) + m_{S_\lambda}^0, \quad (2)$$

where f_λ^0 is the reference spectrum corresponding to a known apparent magnitude $m_{S_\lambda}^0$. The integrands of equation (2) correspond to the total photon energy, which suitably represents data obtained with traditional energy-collecting devices, such as bolometers. However, data from modern photon-collecting devices, such as CCDs and infrared arrays, would be better represented by an integration over the collected photon flux,

$$m_{S_\lambda} = -2.5 \log \left(\frac{\int_{\lambda_1}^{\lambda_2} \lambda f_\lambda S_\lambda d\lambda}{\int_{\lambda_1}^{\lambda_2} \lambda f_\lambda^0 S_\lambda d\lambda} \right) + m_{S_\lambda}^0. \quad (3)$$

The difference between energy integration and photon integration is usually very small when extinction is negligible. In the present paper, we adopt the photon integration (eq. 3) for defining the magnitudes.

By combining the equations above, we derive the absolute magnitude, for which d is defined to be 10 pc:

$$M_{S_\lambda} = -2.5 \log \left[\left(\frac{R}{10\text{pc}} \right)^2 \frac{\int_{\lambda_1}^{\lambda_2} 10^{-0.4A_\lambda} \lambda F_\lambda S_\lambda d\lambda}{\int_{\lambda_1}^{\lambda_2} \lambda f_\lambda^0 S_\lambda d\lambda} \right] + m_{S_\lambda}^0. \quad (4)$$

Stellar radius R can be obtained if the luminosity L and the effective surface temperature T_{eff} of the star are given.

We adopt a Vega-based photometric system (VEGAmag system), which uses Vega (α Lyr) as the calibrating star. Following Girardi et al. (2002), we adopt the synthetic ATLAS9 model (Kurucz 1993) for Vega with $T_{eff} = 9550$ K, $\log g = 3.95$, $[M/H] = -0.5$, microturbulent velocity $\xi = 2$ km/s, $(R/d)^2 = 6.247 \times 10^{-17}$, and photometric zeropoint of 0.03. The dilution factor $(R/d)^2$ is needed to calculate f_λ^{Vega} from the model spectrum F_λ^{Vega} . Now, equation (4) can be written as

$$M_{S_\lambda} = -2.5 \log \left[\frac{(R/10\text{pc})^2 \int_{\lambda_1}^{\lambda_2} 10^{-0.4A_\lambda} \lambda F_\lambda S_\lambda d\lambda}{(R_{Vega}/d_{Vega})^2 \int_{\lambda_1}^{\lambda_2} \lambda F_\lambda^{Vega} S_\lambda d\lambda} \right] + m_{S_\lambda}^{Vega}, \quad (5)$$

where $(R_{Vega}/d_{Vega})^2 = 6.247 \times 10^{-17}$ and $m_{S_\lambda}^{Vega} = 0.03$.

The NICMOS Data Handbook (Dickinson 2002) defines the Vega magnitude system for NICMOS filters by using a filter-averaged flux density of a synthetic model for Vega, $\langle f_\nu^{Vega} \rangle$, as a

photometric zeropoint. Therefore, for NICMOS filters, the magnitude is defined to be

$$M_{S_\lambda} = -2.5 \log \left[\frac{(R/10\text{pc})^2 \int_{\lambda_1}^{\lambda_2} 10^{-0.4A_\lambda} \lambda F_\lambda S_\lambda d\lambda}{\langle f_\nu^{\text{Vega}} \rangle c \int_{\nu_1}^{\nu_2} \nu^{-1} \lambda S_\nu d\nu} \right] + m_{S_\lambda}^{\text{Vega}}. \quad (6)$$

We adopt $\langle f_\nu^{\text{Vega}} \rangle$ values from the NICMOS Data Handbook Ver. 5.0: 1040.7 Jy for F160W, 703.6 Jy for F205W, and 610.4 Jy for F222M.

We adopt filter transmission functions of Bessell & Brett (1988) for H and K , and those of the Quirk camera of the University of Hawaii for K' and K_s .¹ In the case of NICMOS filters, transmission functions include both throughputs and detector quantum efficiencies, which come from SYNPHOT, a synthetic photometry package distributed as a part of STSDAS (Space Telescope Science Data Analysis System) software. When calculating the magnitudes for ground-based filters, we do not consider the atmospheric transmission, telescope throughputs, and detector quantum efficiencies with an assumption that the wavelength dependence of these effects is negligible for the wavelength ranges covered by the filters considered here.

3. Stellar Spectral Library & Evolutionary Tracks

For the spectra of synthetic stellar atmospheres, we adopt Kurucz ATLAS9 no-overshoot models² calculated by Castelli et al. (1997). The metallicities of these models cover the values of $[M/H] = -2.5$ to $+0.5$. A microturbulent velocity $\xi = 2$ km/s, and a mixing length parameter $\alpha = 1.25$ are adopted in the present study.

For the temporal evolution of T_{eff} and L as a function of stellar mass, i.e. stellar evolutionary tracks, we adopt the “basic set” of the Padova models³ (Girardi et al. 2002). We consider isochrones with a metallicity $Z = 0.0001, 0.001, 0.019, \& 0.03$. The stellar spectral library and the evolutionary tracks we adopted assume a solar chemical mixture.

ATLAS9 models cover a wide range of T_{eff} and $\log g$: $3500 \text{ K} \leq T \leq 50000 \text{ K}$, & $0 \leq \log g \leq 5$. However, as shown in Figure 2, some of the $\log T_{\text{eff}}$ and $\log g$ values from the stellar tracks are located outside this range, in particular for small $\log T_{\text{eff}}$ and $\log g$. For this reason, empirical M giant spectra (e.g., Bessell et al. 1989, 1991; Fluks et al. 1994) are often adopted. However, as noted by Girardi et al. (2002), the magnitudes from the ATLAS9 models, and those from the supplementary spectra, do not nicely match at their boundaries. We find that the isochrones in this overlap region are sensitive on how one applies the interpolation between the two libraries. The goal of the present paper is simply to present similarities/disimilarities of the isochrones for

¹<http://www.ifa.hawaii.edu/instrumentaion/quirc/quirc.html>

²“NOVER” files at <http://kurucz.harvard.edu/grids.html>

³<http://pleiadi.pd.astro.it>

different filters, not the most complete isochrones, so we choose not to supplement the ATLAS9 models. Consequently, our isochrones do not include M giants cooler than $T_{eff} \sim 3500$ K, relatively rare metal-rich stars that are usually in the thermally pulsing AGB phase of evolution, and are found at the upper tip of the red giant branch.

We similarly choose not to supplement the ATLAS9 models with synthetic spectra of M, L, & T dwarfs (low T_{eff} and high g), as was done for the Padova models by Girardi et al. (2002) using the spectra calculated by Allard et al. (2000).

4. Isochrones

We first calculate a table of magnitudes for all spectra in ATLAS9 models in the H and K band filters, covering a large range in T_{eff} , $\log g$, and $[M/H]$, using equations (5) and (6). We use this table as a set of interpolates for the T_{eff} , $\log g$, and Z values predicted by the stellar evolution models for a given age in order to estimate synthetic isochrones.

Isochrones for $A_\lambda = 0$, calculated in this way, are shown in Figures 3–6 for four different metallicities, and four ages. As expected for color-magnitude diagrams (CMD) in the infrared, the isochrones have relatively narrow color spreads. The color differences between filters is more prominent for the highest metallicity isochrones. In most cases, isochrones for K' and K_s are nearly indistinguishable, and those for F205W and F222M are quite close to each other. The largest color difference, 0.18 mag, is seen between the bright end of the K and F222M isochrones for $Z = 0.019$ and age = 10^{10} yr. In general, intrinsic color differences of red giants between the atmospheric filters and the space filters are ~ 0.1 -0.15 mag.

Such color differences at the bright end of red giant isochrones between different filters are due to a kink in the continuum near $1.6 \mu\text{m}$ and the CO absorption band longward of $2.3 \mu\text{m}$. Figure 7 compares the spectrum of a K giant with $Z = 0.019$ & Age = 10^9 yr ($K \simeq -5.3$ mag, $T_{eff} = 4000$ K, $\log g = 1.0$), whose F160W–F222M color is 0.13 mag redder than its $H - K$ color, and that of an F dwarf ($K \simeq 0.0$ mag, $T_{eff} = 7000$ K, $\log g = 3.5$), whose F160W–F222M color is only 0.01 mag redder than its $H - K$ color. The spectrum of an F dwarf is very close to that of Vega, and has no strong features in H and K bands, making all colors in our filter sets nearly zero. On the other hand, the spectrum of a K giant 1) significantly deviates from that of Vega at the wavelengths shortward of $1.6 \mu\text{m}$ due to the H^- opacity in cool stars (J. Valenti 2005, private communication), which causes F160W magnitudes to be fainter than H magnitudes, and 2) has the CO absorption band longward of $2.3 \mu\text{m}$, which causes K magnitudes to be fainter than F222M magnitudes. These two effects make F160W–F222M redder than $H - K$, and the same analysis can be applied to other filter sets.

As an independent check of our procedure, we compare our $H - K$ vs. K isochrones to those calculated by Girardi et al. (2002) in Figure 8. The isochrones match nicely, except at the extremes. The discrepancy in the bright end is caused by the empirical M giant spectra that Girardi et al.

(2002) added to their spectral library, and that in the faint end is by the late M dwarf spectra. The discrepancies are considerable only at the top 1 or 2 magnitudes of the isochrone, where only a small number fraction of giants reside.

Magnitude transformations between K -band filters can be obtained from our isochrones. Figures 9 through 13 show magnitude differences between K -band filters as a function of color for all ages considered (10^7 , 10^8 , 10^9 , & 10^{10} yr for $Z = 0.0001$ & 0.019 models, and 6.3×10^7 , 10^8 , 10^9 , & 10^{10} yr for $Z = 0.001$ & 0.03 models), for K -band magnitudes brighter than 4 mag. We find that the magnitude difference can be well fit by a quadratic function for $M < 4$ mag, and by a separate straight line for $M > 4$ mag (we do not present figures for $M > 4$ mag). The largest residuals from the fit are 0.019 mag for the former and 0.009 mag for the latter. The coefficients of the best-fit functions are presented in Tables 1 and 2, along with the residuals and fitting ranges.

Extinction in the K -band is much lower than that in visual bands, but there are still cases where the K -band extinction is significant, such as the central region of the Milky Way ($A_K \sim 3$). Here, we calculate isochrones with K -band extinctions up to 6 mag, some of which are shown in Figures 14–18. For A_λ in equations (5) & (6), we adopt a power-law extinction law for wavelengths between H and K bands:

$$A_\lambda = A_0 \left(\frac{\lambda}{\lambda_0} \right)^{-\alpha}, \quad (7)$$

where we choose $\lambda_0 = 2.2 \mu\text{m}$, and A_0 is the extinction at λ_0 . When assuming that the transmission functions of H and K filters are Dirac delta functions centered at $1.65 \mu\text{m}$ and $2.2 \mu\text{m}$, respectively, the extinction law by Rieke, Rieke, & Paul (1989) gives $\alpha = 1.55$. However, as discussed later in this section, the apparent extinction behaviour of isochrones in the CMD can be different from the actual extinction law, due to non-zero width and asymmetry of the filter transmission functions. We find that $\alpha = 1.61$ makes the isochrone for the $Z = 0.019$, Age = 10^9 yr model behave in the CMD as if it follows an extinction law with $\alpha = 1.55$. We choose this particular isochrone for calibrating the extinction law, with an assumption that the stars used in Rieke et al. (1989) to derive their extinction law, which are the stars in the central parsec of our Galaxy, can be represented by such metallicity and age.

For the sake of easier comparison, isochrones in Figures 14–18 are dereddened by an amount of $A_0(\lambda_c/\lambda_0)^{-1.61}$, where the central wavelength of the filter λ_c is defined by

$$\lambda_c = \frac{\int_{\lambda_1}^{\lambda_2} S_\lambda \lambda d\lambda}{\int_{\lambda_1}^{\lambda_2} S_\lambda d\lambda}, \quad (8)$$

and is given in Table 3. As seen in the figures, isochrones with larger extinction values are more vertically straight, in particular for 10^9 and 10^{10} yr models. This is probably because the extinction makes the longer wavelength part of the spectrum more important, which is closer to the Rayleigh-Jeans regime of the spectrum.

Since we have dereddened the isochrones with the known amount of extinction at λ_c , all the dereddened isochrones with different extinction values in Figures 14–18 should be coincident, if

the filter transmission functions were Dirac delta functions centered at λ_c . The figures show that the dereddened isochrones do not align, and the amount of misalignment is rather large for some filters. This implies that the inferred amount of extinction can be sensitively dependent on the shape of the filter transmission function.

Let us see how this will affect the analyses of actual observations. When one estimates the amount of extinction from an observed near-infrared CMD, where isochrones are nearly vertically straight, one converts an observed color excess to an extinction value following an assumed extinction law, which usually has the form of a power-law. When one has photometric data from a set of two filters, X and Y , the amount of extinction can be estimated by

$$\begin{aligned} A_Y^{est} &= \frac{(m_X - m_Y) - (m_X - m_Y)_0}{A_X/A_Y - 1} \\ &= \frac{(m_X - m_Y) - (m_X - m_Y)_0}{(\lambda_X/\lambda_Y)^{-\alpha} - 1}, \end{aligned} \quad (9)$$

where m_X & m_Y and λ_X & λ_Y are the magnitudes and the central wavelengths of the two filters, respectively, and subscript 0 denotes the intrinsic value. For estimating extinction from our isochrones, we first use $\alpha = 1.55$. Figure 19 shows the difference between the inferred extinction values, using equation (9) and colors from our reddened isochrones, and the actual extinction values. Here, the extinction of each isochrone has been calculated using the mean color (for A_Y^{est}) and magnitude (for A_Y) of the reddened isochrone data points having K -band magnitudes between -6 and 0 mag. As the figure shows, the differences between estimated and actual extinction values are larger for the space filter sets in general. The largest relative difference is $\sim 10\%$, and the largest absolute difference is 0.29 mag. Note that the extinction estimates for the $Z = 0.019$ and Age = 10^9 yr model inferred from H & K are very close to the actual extinction values, justifying our choice of $\alpha = 1.61$ for equation (7).

The problems seen in Figure 19 can be alleviated by finding an “effective extinction slope” for each filter set and isochrone model, which is defined such that it better describes the extinction behaviour in the CMD. Figure 20 shows reddened K -band magnitudes and colors for the $Z = 0.019$ & Age = 10^9 yr isochrone (the figure only shows an isochrone data point whose intrinsic K magnitude is 0, as an example). The reddened magnitudes and colors of each filter set form a nearly straight line (except for filter set F160W & F222M), thus we may assume that A_X/A_Y in equation (9) is not a function of extinction. Then the effective slope of the extinction, α_{eff} , can be calculated by

$$\alpha_{eff} = -\frac{\log(1 + 1/b)}{\log(\lambda_X/\lambda_Y)}, \quad (10)$$

where b is the slope of the straight line that fits the distribution of reddened magnitudes vs. reddened colors, as in Figure 20. We calculate b for data points of each isochrone whose intrinsic K magnitudes are between -6 and 0 mag, and take an average for each isochrone model. When finding the best-fit straight line, we forced the line to include the $A_0 = 0$ point, i.e., we fitted $y = y_0 + b(x - x_0)$, instead of $y = a + bx$, where subscript 0 denotes the values for $A_0 = 0$. Table 4

shows the averages and standard deviations of α_{eff} values for each isochrone model. Note that the standard deviations of α_{eff} in an isochrone is generally much smaller than the differences of average α_{eff} values between different isochrones. The average α_{eff} values range from 1.489 to 1.574, which are 7.5 % to 2.2 % smaller than the original α value we adopted for extinction, 1.61. As seen in Figure 21, extinction values estimated by equation (9) with α_{eff} are now very close to the actual values, except for filter set F160W & F222M, whose extinction behaviour in the CMD is poorly described by a straight line.

The nonlinear extinction behavior of the filter set F160W & F222M appears to be due to a significant width difference in the two filters: F160W covers $\sim 0.4\mu\text{m}$ while F222M covers only $\sim 0.15\mu\text{m}$. Here we discuss the effect of the width difference in terms of an extinction-weighted central wavelength,

$$\lambda_A = \frac{\int_{\lambda_1}^{\lambda_2} 10^{-0.4A_\lambda} S_\lambda \lambda d\lambda}{\int_{\lambda_1}^{\lambda_2} 10^{-0.4A_\lambda} S_\lambda d\lambda}. \quad (11)$$

As the amount of extinction A_0 in equation (7) becomes large, λ_A shifts to a longer wavelength because A_λ decreases with an increasing wavelength. However, the amount of this shift is limited by the filter width. When the widths of two filters differ significantly, the relative shifts in λ_A become considerably different for the two filters. Since the slope of the reddening vector in the CMD is determined by a ratio between λ_A 's of the two filters (see eq. 9), the slope of the reddening vector will become a function of A_0 when the ratio between λ_A 's is sensitively dependent on A_0 , i.e. when the widths of two filters differ significantly. Figure 22 shows the effect of the filter width by comparing the extinction behaviour of five imaginary filter sets. Filter set *a* represents F160W & F222M, and its extinction behaviour in the CMD is not linear, as in the real F160W & F222M set. But this nonlinearity becomes much less significant when the two filters are both wide (filter set *b*) or both narrow (filter set *c*). Filter sets *d* and *e* show that the nonlinearity becomes more prominent when the two filters with significantly different widths are closer to each other. This is because the dependence of λ_A ratio on A_0 is larger when the two filters are closer to each other, and this supports our interpretation of the nonlinear reddening vector of the filter set F160W & F222M.

Now we discuss how one may transform an extinction value estimated for one filter in the K -band to the extinction value for another filter. When the empirical extinction law from observations is not available for the two filters, transformation of extinction values between filters Y and Y' is normally obtained by

$$A_{Y'} = A_Y \left(\frac{\lambda_{Y'}}{\lambda_Y} \right)^{-\alpha}, \quad (12)$$

assuming that the extinction has a functional form of a power-law, and the filter transmission functions are Dirac delta functions centered at their central wavelengths. We find that once A_Y is estimated with α_{eff} , equation (12) with the original α value, 1.61, gives good estimates for $A_{Y'}$. Figure 23 shows the difference between the extinction values that are estimated by equation (9) with our $H - K'$, $H - K_s$, F160W–F205W, & F160W–F222M colors and then converted to an

extinction value at K by equation (12) with $\alpha = 1.61$, and the actual extinction values for K . Except for filter set F160W–F222M again, extinction values can be reliably transformed between different filter sets following the above procedure. It is interesting to note that the original α value of 1.61 is not suitable for estimating the extinction, but can be used for transformation between appropriately estimated extinction values.

5. SUMMARY

We have calculated in a consistent way five near-infrared theoretical isochrones for filter sets $H - K$, $H - K'$, $H - K_s$, F160W–F205W, and F160W–F222M. We presented isochrones for Z of 0.0001 to 0.03, and age of 10^7 to 10^{10} yr. Even in the same Vega magnitude system, near-infrared colors of the same isochrone can be different by up to 0.18 mag at the bright end of the isochrone for different filter sets. The difference of intrinsic colors for a red giant between atmospheric filters and the space filters is generally ~ 0.1 – 0.15 mag. We provided magnitude transformations between K -band filters as a function of color from H & K band filters. We also presented isochrones with A_K of up to 6 mag. Isochrones for larger extinction values are found to be more vertically straight. We found that care is needed when comparing extinction values that are estimated using different filter sets, in particular when comparing those between atmospheric and space filter sets: extinction values inferred using space filters can be in error by up to 0.3 mag. To alleviate this problem, we introduced an “effective extinction slope” for each filter set and isochrone model, which describes the extinction-dependent behaviour of isochrones in the observed CMD.

S.S.K. thanks Hwankyung Sung for helpful discussion. We thank Jeff Valenti for identifying H^- as a source of opacity in cool stars. We thank the anonymous referee whose suggestions and comments significantly improved the content of the present paper. This work was supported by the research fund from Kyung Hee University. S.S.K. was supported by the Astrophysical Research Center for the Structure and Evolution of the Cosmos (ARCSEC) of Korea Science and Engineering Foundation through the Science Research Center (SRC) program. M.G.L. was in part supported by the ABRL (R14-2002-058-01000-0) and the BK21 program.

REFERENCES

- Allard, F., Hauschildt, P. H., Alexander, D. R., Tamanai, A., & Ferguson, J. W. 2000, in Proc. of From giant planets to cool stars, eds. C. A. Griffith, & M. S. Marley, ASP Conf. Ser., 212, 127
- Bessell, M. S., & Brett, J. M. 1988, PASP, 100, 1134
- Bessell, M. S., Brett, J. M., Scholz, M., Wood, P. R. 1989, A&AS, 77, 1
- Bessell, M. S., Brett, J. M., Scholz, M., Wood, P. R. 1991, A&AS, 89, 335
- Castelli, F., Gratton, R. G., & Kurucz, R. L. 1997, A&A, 318, 841
- Dickinson, M., ed. 2002, NICMOS Data Handbook (ver. 5.0; Baltimore: STScI)
- Fluks, M. A., Plez, B., The, P. S., de Winter, D., Westerlund, B. E., & Steenman, H. C. 1994, A&AS, 105, 311
- Girardi, L., Bertelli, G., Bressan, A., Chiosi, C., Groenewegen, M. A. T., Marigo, P., Salsasnich, B., & Weiss, A. 2002, A&A, 391, 195
- Johnson, H. L., Iriarte, B., Mitchell, R. I., & Wisniewskj, W. Z. 1966, Comm. Lunar Plan. Lab., 4, 99
- Kurucz, R. L. 1993, in The Stellar Populations of Galaxies, eds. B. Barbuy, & A. Renzini, (Dordrecht, Kluwer), IAU Symp., 149, 225
- Origlia, L., & Leitherer, C. 2000, AJ, 119, 2018
- Persson, S. E., Murphy, D. C., Krzeminski, W., Roth, M., & Rieke, M. J. 1998, AJ, 116, 2475
- Rieke, G. H., Rieke, M. J., & Paul, A. E. 1989, ApJ, 336, 752
- Stephens, A. W., Grogel, J. A., Ortolani, S., Davies, R., Jablonka, P., Renzini, A., & Rich, R. M. 2000, AJ, 119, 419
- Wainscoat, R. J., & Cowie, L. L. 1992, AJ, 103, 332

Table 1. Best-Fit Coefficients for Magnitude Differences^a ($K < 4$ mag)

Color	Magnitude Difference	Z	c_0	c_1	c_2	Residual ^b (mag)	Fitting Range ^c (mag $\sim$ mag)
$H - K$	$K' - K$	0.0001	-0.000	0.092	-0.166	0.003	-0.108 $\sim$ 0.118
$H - K$	$K' - K$	0.001	0.004	0.018	-3.318	0.009	-0.072 $\sim$ 0.087
$H - K$	$K' - K$	0.019	-0.001	-0.100	-2.285	0.019	-0.096 $\sim$ 0.135
$H - K$	$K' - K$	0.03	-0.003	-0.108	-1.213	0.014	-0.057 $\sim$ 0.168
$H - K$	$K_s - K$	0.0001	0.000	0.039	-0.074	0.003	-0.108 $\sim$ 0.118
$H - K$	$K_s - K$	0.001	0.004	-0.021	-2.580	0.008	-0.072 $\sim$ 0.087
$H - K$	$K_s - K$	0.019	-0.001	-0.130	-1.851	0.016	-0.096 $\sim$ 0.135
$H - K$	$K_s - K$	0.03	-0.002	-0.141	-0.942	0.012	-0.057 $\sim$ 0.168
$H - K$	F205W- K	0.0001	-0.029	0.183	-0.145	0.004	-0.108 $\sim$ 0.118
$H - K$	F205W- K	0.001	-0.026	0.134	-2.389	0.006	-0.072 $\sim$ 0.087
$H - K$	F205W- K	0.019	-0.028	0.060	-1.843	0.013	-0.096 $\sim$ 0.135
$H - K$	F205W- K	0.03	-0.029	0.070	-1.035	0.010	-0.057 $\sim$ 0.168
$H - K$	F222M- K	0.0001	-0.031	-0.042	-0.394	0.005	-0.108 $\sim$ 0.118
$H - K$	F222M- K	0.001	-0.027	-0.113	-3.384	0.008	-0.072 $\sim$ 0.087
$H - K$	F222M- K	0.019	-0.034	-0.240	-2.119	0.018	-0.096 $\sim$ 0.135
$H - K$	F222M- K	0.03	-0.035	-0.271	-0.852	0.014	-0.057 $\sim$ 0.168
$H - K'$	$K - K'$	0.0001	0.000	-0.102	0.228	0.003	-0.096 $\sim$ 0.108
$H - K'$	$K - K'$	0.001	-0.003	-0.057	2.711	0.006	-0.065 $\sim$ 0.110
$H - K'$	$K - K'$	0.019	0.002	0.061	1.356	0.015	-0.086 $\sim$ 0.175
$H - K'$	$K - K'$	0.03	0.002	0.105	0.639	0.011	-0.050 $\sim$ 0.209
$H - K'$	$K_s - K'$	0.0001	0.001	-0.059	0.124	0.003	-0.096 $\sim$ 0.108
$H - K'$	$K_s - K'$	0.001	-0.000	-0.048	0.661	0.002	-0.065 $\sim$ 0.110
$H - K'$	$K_s - K'$	0.019	0.000	-0.034	0.321	0.003	-0.086 $\sim$ 0.175
$H - K'$	$K_s - K'$	0.03	0.000	-0.029	0.184	0.003	-0.050 $\sim$ 0.209
$H - K'$	F205W- K'	0.0001	-0.029	0.100	0.008	0.003	-0.096 $\sim$ 0.108
$H - K'$	F205W- K'	0.001	-0.029	0.111	0.467	0.003	-0.065 $\sim$ 0.110
$H - K'$	F205W- K'	0.019	-0.027	0.147	0.053	0.004	-0.086 $\sim$ 0.175
$H - K'$	F205W- K'	0.03	-0.027	0.165	-0.031	0.004	-0.050 $\sim$ 0.209
$H - K'$	F222M- K'	0.0001	-0.031	-0.147	-0.240	0.005	-0.096 $\sim$ 0.108
$H - K'$	F222M- K'	0.001	-0.032	-0.142	0.409	0.005	-0.065 $\sim$ 0.110
$H - K'$	F222M- K'	0.019	-0.033	-0.140	0.345	0.005	-0.086 $\sim$ 0.175
$H - K'$	F222M- K'	0.03	-0.032	-0.156	0.381	0.005	-0.050 $\sim$ 0.209
$H - K_s$	$K - K_s$	0.0001	-0.000	-0.041	0.086	0.003	-0.104 $\sim$ 0.114
$H - K_s$	$K - K_s$	0.001	-0.003	-0.000	2.027	0.005	-0.070 $\sim$ 0.108
$H - K_s$	$K - K_s$	0.019	0.001	0.095	1.075	0.013	-0.092 $\sim$ 0.174
$H - K_s$	$K - K_s$	0.03	0.002	0.127	0.500	0.009	-0.055 $\sim$ 0.209
$H - K_s$	$K' - K_s$	0.0001	-0.001	0.055	-0.103	0.002	-0.104 $\sim$ 0.114
$H - K_s$	$K' - K_s$	0.001	0.000	0.044	-0.633	0.002	-0.070 $\sim$ 0.108
$H - K_s$	$K' - K_s$	0.019	-0.000	0.032	-0.314	0.003	-0.092 $\sim$ 0.174
$H - K_s$	$K' - K_s$	0.03	-0.000	0.029	-0.184	0.003	-0.055 $\sim$ 0.209
$H - K_s$	F205W- K_s	0.0001	-0.029	0.150	-0.087	0.005	-0.104 $\sim$ 0.114
$H - K_s$	F205W- K_s	0.001	-0.029	0.151	-0.106	0.003	-0.070 $\sim$ 0.108
$H - K_s$	F205W- K_s	0.019	-0.028	0.174	-0.209	0.003	-0.092 $\sim$ 0.174
$H - K_s$	F205W- K_s	0.03	-0.028	0.188	-0.179	0.003	-0.055 $\sim$ 0.209
$H - K_s$	F222M- K_s	0.0001	-0.032	-0.084	-0.335	0.006	-0.104 $\sim$ 0.114

Table 1—Continued

Color	Magnitude Difference	Z	c_0	c_1	c_2	Residual ^b (mag)	Fitting Range ^c (mag $\sim$ mag)
$H - K_s$	F222M– K_s	0.001	–0.032	–0.089	–0.336	0.006	–0.070 $\sim$ 0.108
$H - K_s$	F222M– K_s	0.019	–0.033	–0.100	–0.021	0.007	–0.092 $\sim$ 0.174
$H - K_s$	F222M– K_s	0.03	–0.033	–0.121	0.163	0.007	–0.055 $\sim$ 0.209
F160W–F205W	$K - F205W$	0.0001	0.030	–0.164	0.284	0.007	–0.098 $\sim$ 0.154
F160W–F205W	$K - F205W$	0.001	0.028	–0.168	1.267	0.004	–0.067 $\sim$ 0.169
F160W–F205W	$K - F205W$	0.019	0.030	–0.084	0.681	0.012	–0.087 $\sim$ 0.261
F160W–F205W	$K - F205W$	0.03	0.030	–0.055	0.410	0.009	–0.052 $\sim$ 0.279
F160W–F205W	$K' - F205W$	0.0001	0.029	–0.077	0.064	0.004	–0.098 $\sim$ 0.154
F160W–F205W	$K' - F205W$	0.001	0.029	–0.069	–0.211	0.004	–0.067 $\sim$ 0.169
F160W–F205W	$K' - F205W$	0.019	0.029	–0.111	0.018	0.005	–0.087 $\sim$ 0.261
F160W–F205W	$K' - F205W$	0.03	0.028	–0.117	–0.019	0.004	–0.052 $\sim$ 0.279
F160W–F205W	$K_s - F205W$	0.0001	0.030	–0.126	0.186	0.007	–0.098 $\sim$ 0.154
F160W–F205W	$K_s - F205W$	0.001	0.030	–0.117	0.158	0.005	–0.067 $\sim$ 0.169
F160W–F205W	$K_s - F205W$	0.019	0.029	–0.141	0.185	0.004	–0.087 $\sim$ 0.261
F160W–F205W	$K_s - F205W$	0.03	0.029	–0.141	0.089	0.005	–0.052 $\sim$ 0.279
F160W–F205W	F222M–F205W	0.0001	–0.001	–0.187	0.021	0.005	–0.098 $\sim$ 0.154
F160W–F205W	F222M–F205W	0.001	–0.001	–0.185	0.100	0.002	–0.067 $\sim$ 0.169
F160W–F205W	F222M–F205W	0.019	–0.003	–0.225	0.248	0.005	–0.087 $\sim$ 0.261
F160W–F205W	F222M–F205W	0.03	–0.002	–0.238	0.198	0.003	–0.052 $\sim$ 0.279
F160W–F222M	$K - F222M$	0.0001	0.031	0.020	0.180	0.005	–0.118 $\sim$ 0.188
F160W–F222M	$K - F222M$	0.001	0.029	0.013	0.847	0.005	–0.079 $\sim$ 0.199
F160W–F222M	$K - F222M$	0.019	0.032	0.110	0.358	0.011	–0.105 $\sim$ 0.305
F160W–F222M	$K - F222M$	0.03	0.032	0.143	0.186	0.010	–0.061 $\sim$ 0.333
F160W–F222M	$K' - F222M$	0.0001	0.030	0.092	0.035	0.003	–0.118 $\sim$ 0.188
F160W–F222M	$K' - F222M$	0.001	0.031	0.099	–0.218	0.003	–0.079 $\sim$ 0.199
F160W–F222M	$K' - F222M$	0.019	0.031	0.095	–0.155	0.003	–0.105 $\sim$ 0.305
F160W–F222M	$K' - F222M$	0.03	0.031	0.100	–0.145	0.003	–0.061 $\sim$ 0.333
F160W–F222M	$K_s - F222M$	0.0001	0.031	0.051	0.114	0.005	–0.118 $\sim$ 0.188
F160W–F222M	$K_s - F222M$	0.001	0.031	0.058	0.047	0.005	–0.079 $\sim$ 0.199
F160W–F222M	$K_s - F222M$	0.019	0.032	0.069	–0.032	0.005	–0.105 $\sim$ 0.305
F160W–F222M	$K_s - F222M$	0.03	0.031	0.079	–0.067	0.005	–0.061 $\sim$ 0.333
F160W–F222M	F205W–F222M	0.0001	0.000	0.157	–0.003	0.004	–0.118 $\sim$ 0.188
F160W–F222M	F205W–F222M	0.001	0.001	0.157	–0.061	0.002	–0.079 $\sim$ 0.199
F160W–F222M	F205W–F222M	0.019	0.002	0.185	–0.150	0.005	–0.105 $\sim$ 0.305
F160W–F222M	F205W–F222M	0.03	0.002	0.194	–0.115	0.003	–0.061 $\sim$ 0.333

Note. — Only the data points that have $\log T_{eff} \geq 3500$ K and $\log g \geq 0$ were considered for the fitting.

^aMagnitude differences are fitted to a function $[\text{Mag Diff}] = c_0 + c_1[\text{Color}] + c_2[\text{Color}]^2$.

^bThe largest absolute residual.

^cColor range where the fit is valid.

Table 2. Best-Fit Coefficients for Magnitude Differences^a ($K > 4$ mag)

Color	Magnitude Difference	Z	c_0	c_1	Residual ^b (mag)	Fitting Range ^c (mag $\sim$ mag)
$H - K$	$K' - K$	0.0001	0.001	0.112	0.001	0.026 $\sim$ 0.195
$H - K$	$K' - K$	0.001	-0.001	0.097	0.003	0.028 $\sim$ 0.223
$H - K$	$K' - K$	0.019	-0.014	0.131	0.004	0.046 $\sim$ 0.242
$H - K$	$K' - K$	0.03	-0.017	0.136	0.005	0.050 $\sim$ 0.235
$H - K$	$K_s - K$	0.0001	0.002	0.053	0.001	0.026 $\sim$ 0.195
$H - K$	$K_s - K$	0.001	0.001	0.029	0.002	0.028 $\sim$ 0.223
$H - K$	$K_s - K$	0.019	-0.011	0.060	0.003	0.046 $\sim$ 0.242
$H - K$	$K_s - K$	0.03	-0.013	0.060	0.004	0.050 $\sim$ 0.235
$H - K$	F205W- K	0.0001	-0.027	0.210	0.001	0.026 $\sim$ 0.195
$H - K$	F205W- K	0.001	-0.030	0.198	0.003	0.028 $\sim$ 0.223
$H - K$	F205W- K	0.019	-0.041	0.232	0.004	0.046 $\sim$ 0.242
$H - K$	F205W- K	0.03	-0.042	0.233	0.004	0.050 $\sim$ 0.235
$H - K$	F222M- K	0.0001	-0.032	-0.050	0.001	0.026 $\sim$ 0.195
$H - K$	F222M- K	0.001	-0.033	-0.066	0.003	0.028 $\sim$ 0.223
$H - K$	F222M- K	0.019	-0.046	-0.037	0.003	0.046 $\sim$ 0.242
$H - K$	F222M- K	0.03	-0.048	-0.038	0.003	0.050 $\sim$ 0.235
$H - K'$	$K - K'$	0.0001	-0.001	-0.126	0.001	0.022 $\sim$ 0.172
$H - K'$	$K - K'$	0.001	0.001	-0.106	0.003	0.024 $\sim$ 0.201
$H - K'$	$K - K'$	0.019	0.017	-0.149	0.005	0.050 $\sim$ 0.225
$H - K'$	$K - K'$	0.03	0.020	-0.160	0.004	0.058 $\sim$ 0.219
$H - K'$	$K_s - K'$	0.0001	0.000	-0.066	0.001	0.022 $\sim$ 0.172
$H - K'$	$K_s - K'$	0.001	0.002	-0.074	0.002	0.024 $\sim$ 0.201
$H - K'$	$K_s - K'$	0.019	0.004	-0.081	0.001	0.050 $\sim$ 0.225
$H - K'$	$K_s - K'$	0.03	0.005	-0.089	0.001	0.058 $\sim$ 0.219
$H - K'$	F205W- K'	0.0001	-0.028	0.110	0.001	0.022 $\sim$ 0.172
$H - K'$	F205W- K'	0.001	-0.029	0.111	0.001	0.024 $\sim$ 0.201
$H - K'$	F205W- K'	0.019	-0.028	0.116	0.001	0.050 $\sim$ 0.225
$H - K'$	F205W- K'	0.03	-0.027	0.112	0.001	0.058 $\sim$ 0.219
$H - K'$	F222M- K'	0.0001	-0.033	-0.182	0.001	0.022 $\sim$ 0.172
$H - K'$	F222M- K'	0.001	-0.032	-0.180	0.002	0.024 $\sim$ 0.201
$H - K'$	F222M- K'	0.019	-0.028	-0.193	0.002	0.050 $\sim$ 0.225
$H - K'$	F222M- K'	0.03	-0.028	-0.203	0.002	0.058 $\sim$ 0.219
$H - K_s$	$K - K_s$	0.0001	-0.002	-0.056	0.001	0.024 $\sim$ 0.184
$H - K_s$	$K - K_s$	0.001	-0.001	-0.030	0.002	0.025 $\sim$ 0.215
$H - K_s$	$K - K_s$	0.019	0.012	-0.063	0.004	0.051 $\sim$ 0.239
$H - K_s$	$K - K_s$	0.03	0.014	-0.065	0.004	0.059 $\sim$ 0.234
$H - K_s$	$K' - K_s$	0.0001	-0.000	0.062	0.001	0.024 $\sim$ 0.184
$H - K_s$	$K' - K_s$	0.001	-0.002	0.069	0.002	0.025 $\sim$ 0.215
$H - K_s$	$K' - K_s$	0.019	-0.004	0.075	0.001	0.051 $\sim$ 0.239
$H - K_s$	$K' - K_s$	0.03	-0.005	0.082	0.001	0.059 $\sim$ 0.234
$H - K_s$	F205W- K_s	0.0001	-0.028	0.166	0.001	0.024 $\sim$ 0.184
$H - K_s$	F205W- K_s	0.001	-0.030	0.173	0.002	0.025 $\sim$ 0.215
$H - K_s$	F205W- K_s	0.019	-0.032	0.182	0.001	0.051 $\sim$ 0.239
$H - K_s$	F205W- K_s	0.03	-0.031	0.184	0.001	0.059 $\sim$ 0.234
$H - K_s$	F222M- K_s	0.0001	-0.034	-0.109	0.001	0.024 $\sim$ 0.184

Table 2—Continued

Color	Magnitude Difference	Z	c_0	c_1	Residual ^b (mag)	Fitting Range ^c (mag $\sim$ mag)
$H - K_s$	F222M– K_s	0.001	–0.034	–0.098	0.001	0.025 $\sim$ 0.215
$H - K_s$	F222M– K_s	0.019	–0.033	–0.103	0.001	0.051 $\sim$ 0.239
$H - K_s$	F222M– K_s	0.03	–0.034	–0.105	0.001	0.059 $\sim$ 0.234
F160W–F205W	$K - F205W$	0.0001	0.032	–0.217	0.001	0.047 $\sim$ 0.211
F160W–F205W	$K - F205W$	0.001	0.036	–0.194	0.005	0.054 $\sim$ 0.246
F160W–F205W	$K - F205W$	0.019	0.059	–0.269	0.009	0.092 $\sim$ 0.274
F160W–F205W	$K - F205W$	0.03	0.063	–0.277	0.009	0.102 $\sim$ 0.269
F160W–F205W	$K' - F205W$	0.0001	0.030	–0.101	0.001	0.047 $\sim$ 0.211
F160W–F205W	$K' - F205W$	0.001	0.032	–0.100	0.003	0.054 $\sim$ 0.246
F160W–F205W	$K' - F205W$	0.019	0.035	–0.118	0.002	0.092 $\sim$ 0.274
F160W–F205W	$K' - F205W$	0.03	0.034	–0.114	0.002	0.102 $\sim$ 0.269
F160W–F205W	$K_s - F205W$	0.0001	0.032	–0.163	0.001	0.047 $\sim$ 0.211
F160W–F205W	$K_s - F205W$	0.001	0.036	–0.166	0.004	0.054 $\sim$ 0.246
F160W–F205W	$K_s - F205W$	0.019	0.044	–0.200	0.004	0.092 $\sim$ 0.274
F160W–F205W	$K_s - F205W$	0.03	0.044	–0.206	0.004	0.102 $\sim$ 0.269
F160W–F205W	F222M–F205W	0.0001	0.001	–0.269	0.001	0.047 $\sim$ 0.211
F160W–F205W	F222M–F205W	0.001	0.005	–0.260	0.005	0.054 $\sim$ 0.246
F160W–F205W	F222M–F205W	0.019	0.017	–0.314	0.006	0.092 $\sim$ 0.274
F160W–F205W	F222M–F205W	0.03	0.018	–0.322	0.007	0.102 $\sim$ 0.269
F160W–F222M	$K - F222M$	0.0001	0.031	0.041	0.001	0.058 $\sim$ 0.267
F160W–F222M	$K - F222M$	0.001	0.031	0.052	0.002	0.066 $\sim$ 0.309
F160W–F222M	$K - F222M$	0.019	0.043	0.034	0.003	0.110 $\sim$ 0.344
F160W–F222M	$K - F222M$	0.03	0.046	0.034	0.002	0.124 $\sim$ 0.339
F160W–F222M	$K' - F222M$	0.0001	0.029	0.132	0.001	0.058 $\sim$ 0.267
F160W–F222M	$K' - F222M$	0.001	0.028	0.128	0.003	0.066 $\sim$ 0.309
F160W–F222M	$K' - F222M$	0.019	0.020	0.150	0.004	0.110 $\sim$ 0.344
F160W–F222M	$K' - F222M$	0.03	0.019	0.158	0.004	0.124 $\sim$ 0.339
F160W–F222M	$K_s - F222M$	0.0001	0.031	0.084	0.001	0.058 $\sim$ 0.267
F160W–F222M	$K_s - F222M$	0.001	0.031	0.075	0.001	0.066 $\sim$ 0.309
F160W–F222M	$K_s - F222M$	0.019	0.028	0.087	0.002	0.110 $\sim$ 0.344
F160W–F222M	$K_s - F222M$	0.03	0.028	0.089	0.002	0.124 $\sim$ 0.339
F160W–F222M	F205W–F222M	0.0001	–0.001	0.212	0.001	0.058 $\sim$ 0.267
F160W–F222M	F205W–F222M	0.001	–0.004	0.208	0.004	0.066 $\sim$ 0.309
F160W–F222M	F205W–F222M	0.019	–0.013	0.240	0.005	0.110 $\sim$ 0.344
F160W–F222M	F205W–F222M	0.03	–0.014	0.245	0.005	0.124 $\sim$ 0.339

Note. — Only the data points that have $\log T_{eff} \geq 3500$ K and $\log g \geq 0$ were considered for the fitting.

^aMagnitude differences are fitted to a function $[\text{Mag Diff}] = c_0 + c_1[\text{Color}]$.

^bThe largest absolute residual.

^cColor range where the fit is valid.

Table 3. Central Wavelength λ_c (μm)

H	K	K'	K_s	F160W	F205W	F222M
1.646	2.212	2.114	2.160	1.610	2.079	2.219

Note. — λ_c is defined by equation (8).

Table 4. Averages & Standard Deviations of α_{eff} Values

Isochrone Model		$H - K$	$H - K'$	$H - K_s$	F160W–F205W	F160W–F222M
Z	Age					
0.0001	10^7	1.574 ± 0.001	1.572 ± 0.001	1.574 ± 0.001	1.544 ± 0.001	1.566 ± 0.001
0.0001	10^8	1.570 ± 0.002	1.567 ± 0.003	1.569 ± 0.003	1.539 ± 0.005	1.557 ± 0.006
0.0001	10^9	1.566 ± 0.002	1.563 ± 0.002	1.565 ± 0.002	1.528 ± 0.008	1.544 ± 0.008
0.0001	10^{10}	1.564 ± 0.002	1.560 ± 0.002	1.562 ± 0.002	1.520 ± 0.007	1.537 ± 0.006
0.001	6.3×10^7	1.570 ± 0.003	1.567 ± 0.003	1.570 ± 0.003	1.539 ± 0.005	1.557 ± 0.007
0.001	10^8	1.568 ± 0.004	1.565 ± 0.003	1.568 ± 0.004	1.536 ± 0.009	1.553 ± 0.009
0.001	10^9	1.562 ± 0.005	1.560 ± 0.003	1.563 ± 0.003	1.520 ± 0.009	1.538 ± 0.006
0.001	10^{10}	1.556 ± 0.009	1.557 ± 0.005	1.559 ± 0.005	1.511 ± 0.013	1.532 ± 0.009
0.019	10^7	1.574 ± 0.001	1.571 ± 0.001	1.574 ± 0.001	1.545 ± 0.001	1.566 ± 0.001
0.019	10^8	1.561 ± 0.013	1.562 ± 0.008	1.564 ± 0.009	1.526 ± 0.021	1.547 ± 0.017
0.019	10^9	1.549 ± 0.009	1.555 ± 0.006	1.555 ± 0.006	1.505 ± 0.015	1.529 ± 0.010
0.019	10^{10}	1.539 ± 0.010	1.548 ± 0.007	1.548 ± 0.007	1.489 ± 0.016	1.517 ± 0.012
0.03	6.3×10^7	1.568 ± 0.008	1.567 ± 0.005	1.569 ± 0.006	1.538 ± 0.012	1.557 ± 0.011
0.03	10^8	1.560 ± 0.014	1.562 ± 0.008	1.563 ± 0.009	1.526 ± 0.022	1.547 ± 0.017
0.03	10^9	1.547 ± 0.010	1.554 ± 0.006	1.554 ± 0.006	1.503 ± 0.014	1.528 ± 0.010
0.03	10^{10}	1.538 ± 0.009	1.548 ± 0.006	1.548 ± 0.006	1.489 ± 0.013	1.517 ± 0.010

Note. — Data are presented in the form of [Average $\pm$ Standard Deviation]. The average and standard deviation values are calculated from the data points of each isochrone whose intrinsic K magnitudes are between -6 and 0 mag.




Fig. 1.— Transmission functions (S_λ) of the filters considered in the present work. S_λ 's are scaled such that their maximum values become 1.




Fig. 2.— $[\log T_{eff}, \log g]$ distribution of the spectra that we use in the present work among the stellar spectral library by Castelli et al. (1997; crosses), and the same distribution of the stellar models that we use among the stellar evolutionary tracks by Girardi et al. (2002; dots).




Fig. 3.— Isochrones of $Z = 0.0001$ model for $H - K$ vs. K (solid), $H - K'$ vs. K' (long dash), $H - K_s$ vs. K_s (short dash), F160W–F205W vs. F205W (long dash-dot), and F160W–F222M vs. F222M (short dash-dot), in Vega magnitude system. The three atmospheric filters nearly coincide at the bright end. Only the data points that have $\log T_{eff} \geq 3500$ K and $\log g \geq 0$ are plotted.




Fig. 4.— Same as Figure 3 but for $Z = 0.001$ model. Isochrones for K' and K_s are indistinguishable.




Fig. 5.— Same as Figure 3 but for $Z = 0.019$ model. Isochrones for K' and K_s are indistinguishable.




Fig. 6.— Same as Figure 3 but for $Z = 0.03$ model. Isochrones for K' and K_s are indistinguishable.




Fig. 7.— ATLAS9 stellar spectra for a K giant (thick solid lines; $T_{eff} = 4000 \text{ K}$, $\log g = 1.0$), an F dwarf (dashed lines; $T_{eff} = 7000 \text{ K}$, $\log g = 3.5$), and Vega (dotted lines). Also plotted in panels *b* and *c* are the transmission functions (S_λ) of *H*, *K*, *F160W*, and *F222M* filters (thin solid lines).




Fig. 8.— $H - K$ vs. K isochrones calculated in the present study (solid lines) and those by Girardi et al. (2002; dashed lines). Only the data points that have $\log T_{eff} \geq 3500$ K and $\log g \geq 0$ are plotted.




Fig. 9.— Magnitude differences between K and the other K -band filters as a function of $H - K$ color for all ages considered in the present paper (10^7 , 10^8 , 10^9 , & 10^{10} yr for $Z = 0.0001$ & 0.019 models, and 6.3×10^7 , 10^8 , 10^9 , & 10^{10} yr for $Z = 0.001$ & 0.03 models), for K -band magnitudes brighter than 4 mag. Best-fit 2nd-order polynomials are also shown. Only the data points that have $\log T_{eff} \geq 3500$ K and $\log g \geq 0$ are plotted.




Fig. 10.— Same as Figure 9 but for magnitude differences between K' and the other K -band filters as a function of $H - K'$ color.




Fig. 11.— Same as Figure 9 but for magnitude differences between K_s and the other K -band filters as a function of $H - K_s$ color.




Fig. 12.— Same as Figure 9 but for magnitude differences between F205W and the other K -band filters as a function of $F160W - F205W$ color.




Fig. 13.— Same as Figure 9 but for magnitude differences between F222M and the other K -band filters as a function of $F160W-F222M$ color.




Fig. 14.— Dereddened $H - K$ vs. K isochrones of $Z = 0.019$ model with $A_0=0$ (solid), $A_0=2$ (long dash), $A_0=4$ (short dash), and $A_0=6$ (dots). The isochrones are dereddened by an amount of $A_0(\lambda_c/\lambda_0)^{-1.61}$. Only the data points that have $\log T_{eff} \geq 3500$ K and $\log g \geq 0$ are plotted.




Fig. 15.— Same as Figure 14 but for $H - K'$ vs. K' .




Fig. 16.— Same as Figure 14 but for $H - K_s$ vs. K_s .




Fig. 17.— Same as Figure 14 but for F160W-F205W vs. F205W.




Fig. 18.— Same as Figure 14 but for $F160W-F222M$ vs. $F222M$.




Fig. 19.— The difference between the extinction values that are estimated by equation (9) with the colors from our reddened isochrones and the actual extinction values. A constant value of 1.55 is used for α in equation (9). The extinction of each isochrone has been estimated with the mean color (for A_Y^{est}) and the mean magnitude (for A_Y) of the reddened isochrone data points whose intrinsic K -band magnitudes are between -6 and 0 mag.




Fig. 20.— Reddened magnitudes of K -band filters and reddened colors for the $Z = 0.019$ & Age = 10^9 yr isochrone data point whose intrinsic K magnitude is 0. Also shown are the best-fit straight lines that go through the data point for $A_\lambda = 0$ for each filter set.




Fig. 21.— Same as Figure 19 but using α_{eff} for equation (9).




Fig. 22.— Reddened magnitudes and colors (crosses) for the $Z = 0.019$ & Age = 10^9 yr isochrone data point whose intrinsic K magnitude is 0, for five imaginary filter sets whose transmission functions are shown in the right panel. Also shown are the best-fit straight lines that go through the data point for $A_\lambda = 0$ for each filter set.




Fig. 23.— The difference between the extinction values that are estimated by equation (9) with our $H - K'$, $H - K_s$, $F160W - F205W$, & $F160W - F222M$ colors and then converted to an extinction value at K by equation (12) with $\alpha = 1.61$, and the extinction values for K that are estimated by equation (9) with our $H - K$ colors.